arXiv:0707.4020v2 [quant-ph] 9 Sep 2007

Geometric measures of entanglement and the Schmidt decomposition

D. Ostapchuk,[*] G. Passante,[†] R. Kobes,[‡] and G. Kunstatter[§]

Physics Department
and
Winnipeg Institute for Theoretical Physics
University of Winnipeg
Winnipeg, MB R3B 2E9
Canada

In the standard geometric approach to a measure of entanglement of a pure state, $\sin^2\theta$ is used, where θ is the angle between the state to the closest separable state of products of normalized qubit states. We consider here a generalization of this notion to separable states consisting of products of unnormalized states of different dimension. In so doing, the entanglement measure $\sin^2\theta$ is found to have an interpretation as the distance between the state to the closest separable state. We also find the components of the closest separable state and its norm have an interpretation in terms of, respectively, the eigenvectors and eigenvalues of the reduced density matrices arising in the Schmidt decomposition of the state vector.

PACS numbers: 03.65.Ud, 03.67.Mn

I. INTRODUCTION

With recognition of its role as a resource in quantum computing [1], the nature of entanglement in quantum systems is a problem of much current interest [2, 3, 4]. Of particular importance is a quantitative measure of entanglement [5]; two of the more commonly used measures, depending on the context, are the von Neumann entropy, based on reduced density matrices [2], and a geometric measure, based on the distance to the nearest product state [6, 7, 8, 9, 10].

The use of the von Neumann entropy as a measure of entanglement can be understood in the framework of the Schmidt decomposition [1, 11, 12, 13, 14, 15]. Consider a pure state $|\psi\rangle$ in an n-dimensional Hilbert space, assumed to be normalized: $\langle\psi|\psi\rangle = 1$. Let us then decompose the system into an u-dimensional subsystem, A, and a v-dimensional subsystem, B, such that $n = uv$. For a basis $|i\rangle$ of A and $|j\rangle$ of B we can write $|\psi\rangle$ as

$$|\psi\rangle = \sum_{i=0}^{u-1}\sum_{j=0}^{v-1}\gamma_{ij}|i\rangle\otimes|j\rangle \tag{1}$$

for some complex coefficients γ_{ij}. However, the Schmidt decomposition states that there exists a basis $|\alpha_i\rangle$ for A and $|\beta_j\rangle$ for B such that $|\psi\rangle$ can be expressed as a single summation:

$$|\psi\rangle = \sum_{k=0}^{\min(u-1,v-1)} \sqrt{p_k}\ |\alpha_k\rangle\otimes|\beta_k\rangle \tag{2}$$

where the Schmidt coefficients satisfy $\sum_k p_k = 1$. The existence of this decomposition follows from the singular value decomposition of the matrix of coefficients $\gamma_{ij}|i\rangle\otimes\langle j|$, and can be related to the reduced density matrices formed by tracing out one of the subsystems of the density matrix $\rho = |\psi\rangle\langle\psi|$:

$$\begin{aligned}\rho_A &= \mathrm{Tr}_B(\rho) = \sum_k p_k\ |\alpha_k\rangle\langle\alpha_k| \\ \rho_B &= \mathrm{Tr}_A(\rho) = \sum_k p_k\ |\beta_k\rangle\langle\beta_k| \end{aligned}\tag{3}$$

---

[*] Electronic address: dcmostap@uwaterloo.ca
[†] Electronic address: glpassan@mail.iqc.uwaterloo.ca
[‡] Electronic address: r.kobes@uwinnipeg.ca
[§] Electronic address: g.kunstatter@uwinnipeg.ca

by which one can see that α_k are the eigenvectors of $\rho_A = \mathrm{Tr}_B(\rho)$ and β_k are the eigenvectors of $\rho_B = \mathrm{Tr}_A(\rho)$. The spectrum of eigenvalues of ρ_A and ρ_B, which are the same, can be used to quantify the degree of entanglement of $|\psi\rangle$ due to the fact that, for a separable state, only one non–zero eigenvalue is present. In this context $K = 1/\sum_k p_k^2$, which satisfies $K \geq 1$, is often considered; another commonly used measure is the von Neumann entropy:

$$S = -\mathrm{Tr}\left(\rho_A \ln \rho_A\right) = -\mathrm{Tr}\left(\rho_B \ln \rho_B\right) = -\sum_k p_k \log_2 p_k \tag{4}$$

On the other hand, in the geometric approach to measuring entanglement, one considers the distance from a n–dimensional pure state $|\psi\rangle = \sum_{i=0}^{n-1} \chi_{p_i} |e_{p_i}^{(i)}\rangle$ to a separable state

$$|\phi\rangle = \otimes_{i=0}^{n-1} |\phi^{(i)}\rangle = \sum_{p_i} c_{p_i}^{(i)} |e_{p_i}^{(i)}\rangle \tag{5}$$

where $i = 0, 1, \ldots, n-1$. The states $|\phi^{(i)}\rangle$ here are assumed to be normalized: $\langle \phi^{(i)} | \phi^{(i)} \rangle = 1$. Minimizing $||\phi\rangle - |\psi\rangle|^2$ results in the non–linear eigenvalue equations:

$$\sum_{p_0 \cdots \widehat{p_i} \cdots p_{n-1}} \chi^*_{p_0 \cdots p_{n-1}} c_{p_0}^{(0)} \cdots \widehat{c_{p_i}^{(i)}} \cdots c_{p_{n-1}}^{(n-1)} = \Lambda c_{p_i}^{(i)*} \tag{6}$$

where the eigenvalue Λ is the Lagrange multiplier enforcing $\langle \phi^{(i)} | \phi^{(i)} \rangle = 1$ and $\widehat{\ }$ means omission. The eigenvalues Λ can be shown to lie in the range $-1 \leq \Lambda \leq 1$, and so are interpreted as the cosine of the angle between $|\phi\rangle$ and $|\psi\rangle$; this can be seen by multiplying Eq. (6) by $c_{p_i}^{(i)}$ and summing over i. A measure of entanglement is then taken to be $1 - \Lambda_{\max}^2$, where $\Lambda_{\max}$ corresponds to the eigenvalue determined from Eq. (6) of the closest separable state.

In this paper we show that if one uses *unnormalized* separable states in a geometric measure of entanglement, the norm of the closest separable state can be related to both the distance between and to the angle between the separable and target states; this will provide a natural interpretation of the entanglement measure $1 - \Lambda_{\max}^2$ as the distance to the closest separable state. As well, by considering the geometric measure in arbitrary dimensional spaces, a connection can be established between the basis states and eigenvalues of the Schmidt decomposition of the state and the components and norm of the closest separable product state used in this geometric approach.

II. OPTIMUM EUCLIDEAN DISTANCE

In this section we describe the geometric measure of entanglement we shall use based on finding the extrema of the distance to a separable state. For this, we consider an n–dimensional pure state $|\psi\rangle = \sum_{i=0}^{n-1} \chi_{p_i} |e_{p_i}^{(i)}\rangle$. We split the n–dimensional space into subspaces $A, B, C, \ldots$ of dimension $u, v, \ldots$, and consider the separable state

$$|\phi\rangle = |A\rangle \otimes |B\rangle \otimes |C\rangle \otimes \ldots = \sum_{i=0}^{u-1} a_i |e_A^{(i)}\rangle \otimes \sum_{j=0}^{v-1} b_j |e_B^{(j)}\rangle \otimes \sum_{k=0}^{w-1} c_j |e_C^{(k)}\rangle \otimes \ldots \tag{7}$$

The state $|\phi\rangle$ is not assumed to be normalized. We now form the distance from the state $|\psi\rangle$ to such a separable state:

$$D^2 = ||\phi\rangle - |\psi\rangle|^2 = \left(\sum_{i=0}^{u-1} \sum_{j=0}^{v-1} \sum_{k=0}^{w-1} \cdots \right) \left(a_i^* b_j^* c_k^* \ldots - \chi^*_{ijk\cdots} \right) \left(a_i b_j c_k \ldots - \chi_{ijk\cdots} \right) \tag{8}$$

where $a_i, b_j, \ldots$ are the coordinates in the appropriate spaces, and optimize this distance with respect to the coordinates of $|A\rangle, |B\rangle, \ldots$:

$$\begin{aligned}
\frac{\partial D^2}{\partial a_i} = 0 \;&\Rightarrow\; a_i^* N_B N_C \ldots = \left(\sum_{j=0}^{v-1} \sum_{k=0}^{w-1} \cdots \right) b_j c_k \ldots \chi^*_{ijk\ldots} \\
\frac{\partial D^2}{\partial b_j} = 0 \;&\Rightarrow\; b_j^* N_A N_C \ldots = \left(\sum_{i=0}^{u-1} \sum_{k=0}^{w-1} \cdots \right) a_i c_k \ldots \chi^*_{ijk\ldots} \\
\frac{\partial D^2}{\partial c_k} = 0 \;&\Rightarrow\; c_k^* N_A N_B \ldots = \left(\sum_{i=0}^{u-1} \sum_{j=0}^{v-1} \cdots \right) a_i b_j \ldots \chi^*_{ijk\ldots} \\
&\vdots
\end{aligned} \tag{9}$$

where $N_A = \sum_{i=0}^{u-1} a_i^* a_i$, $N_B = \sum_{j=0}^{v-1} b_j^* b_j$, and so on. Except in special cases these non–linear equations must be solved numerically. However, from Eq.(9), one can show

$$N_A N_B N_C \ldots = \left(\sum_{i=0}^{u-1} \sum_{j=0}^{v-1} \sum_{k=0}^{w-1} \cdots \right) a_i b_j c_k \ldots \chi^*_{ijk\ldots} \tag{10}$$

and hence, at the critical points,

$$\langle \phi | \psi \rangle = N_A N_B N_C \ldots = \langle \psi | \phi \rangle \tag{11}$$

Using $\langle \psi | \psi \rangle = 1$ and $\langle \phi | \phi \rangle = N_A N_B N_C \ldots$, we find that, at the critical point, the angle between $|\phi\rangle$ and $|\psi\rangle$ is

$$\cos \theta_C = \left. \frac{\langle \psi | \phi \rangle}{\sqrt{\langle \phi | \phi \rangle} \sqrt{\langle \psi | \psi \rangle}} \right|_{\text{critical}} = \sqrt{N_A N_B N_C \ldots} \tag{12}$$

and the distance D^2 is

$$D_C^2 = 1 - N_A N_B N_C \ldots = \sin^2 \theta_C \tag{13}$$

Consistency of the above requires $N_A N_B N_C \ldots \leq 1$. This can be done using the Cauchy–Schwartz inequality:

$$|\langle \phi - \psi | \phi \rangle|^2 \leq \langle \phi - \psi | \phi - \psi \rangle \langle \phi | \phi \rangle \tag{14}$$

Using $\langle \phi | \phi \rangle = N_A N_B N_C \cdots$ and the definition $D^2 = \langle \phi - \psi | \phi - \psi \rangle$, this then implies

$$N_A N_B N_C \cdots D^2 \geq |\langle \phi | \phi \rangle - \langle \psi | \phi \rangle|^2 \tag{15}$$

At the extremal points we have $\langle \psi | \phi \rangle = N_A N_B N_C \cdots$ and $D^2 = 1 - N_A N_B N_C \cdots$, and so we can conclude

$$1 - N_A N_B N_C \cdots \geq 0 \Rightarrow N_A N_B N_C \cdots \leq 1 \tag{16}$$

The preceding has a close connection to the results of the entanglement measure using a normalized separable state; indeed, rewriting Eqs. (9) for the unnormalized separable state in terms of variables $\widetilde{a}_i = a_i / \sqrt{N_A}$, $\widetilde{b}_j = b_j / \sqrt{N_B}$, etc. formally leads to the relations of Eq. (6), with the eigenvalue Λ identified with $\sqrt{\langle \phi | \phi \rangle} = \sqrt{N_A N_B N_C \ldots}$. This makes for a simple geometrical interpretation of the results of using unnormalized and normalized separable states indicated in Fig. (1). Both the approach of Eq. (6) using normalized separable states and that of Eq. (9) using unnormalized states lead to the same angle $\cos \theta_C = \Lambda = \sqrt{N_A N_B N_C \ldots}$ of Eq. (12). The corresponding distances differ, however. Using unnormalized separable states $|\phi\rangle$, the distance of Eq. (13) is

$$D^2 = \langle \phi - \psi | \phi - \psi \rangle = 1 - \left(\sqrt{\langle \phi | \phi \rangle} \right)^2 = 1 - \cos^2 \theta_C, \tag{17}$$

On the other hand, using normalized separable states $|\phi_N\rangle$, the corresponding distance is

$$D_N^2 = \langle \phi_N - \psi | \phi_N - \psi \rangle = D^2 + \left(1 - \sqrt{\langle \phi | \phi \rangle} \right)^2 = 2(1 - \cos \theta_C). \tag{18}$$

Thus, in the approach using unnormalized separable states, the use of $1 - \cos \theta_C^2 = \sin \theta_C^2$ as the entanglement measure can be interpreted as the distance to the closest separable state.

III. SCHMIDT DECOMPOSITION

As mentioned in the previous section, the equations of (9) determining the extremal points of the distance to the closest separable state are non–linear. One of the special cases for which a closed–form solution exists is when the


$\vec{\psi}$
$\sqrt{\langle\psi|\psi\rangle}=1$
D
D_N
θ_C
$\sqrt{\langle\phi|\phi\rangle}$
$1-\sqrt{\langle\phi|\phi\rangle}$
$\vec{\phi}$
$\vec{\phi}_{\mathbf{N}}$
$\sqrt{\langle\phi_N|\phi_N\rangle}=1$

FIG. 1: Geometrical comparison of the entanglement measure using unnormalized ($\vec{\phi}$) and normalized ($\vec{\phi}_N$) separable states

n–dimensional system is decomposed into an u-dimensional subsystem, A, and a v-dimensional subsystem, B, such that $n = uv$. In this case the equations decouple to yield

$$\begin{aligned} a_k N_A N_B &= \sum_{i=0}^{u-1}\sum_{j=0}^{v-1} \chi_{kj}\chi_{ij}^* a_i \\ b_m N_A N_B &= \sum_{j=0}^{v-1}\sum_{i=0}^{u-1} \chi_{im}\chi_{ij}^* b_j \end{aligned} \tag{19}$$

which, respectively, leads to the product $N_A N_B$ being found as solutions to

$$\begin{aligned} \det\left| N_A N_B \delta_{ki} - \sum_{j=0}^{v-1} \chi_{kj}\chi_{ij}^* \right| &= 0 \\ \det\left| N_A N_B \delta_{mj} - \sum_{i=0}^{u-1} \chi_{im}\chi_{ij}^* \right| &= 0 \end{aligned} \tag{20}$$

This has an interesting correspondence to the Schmidt decomposition of $|\psi\rangle$ of Eq. (2). To see this, consider the density matrix of the pure state $|\psi\rangle$:

$$\rho = |\psi\rangle\langle\psi| = \sum_{i=0}^{n-1}\sum_{j=0}^{n-1} \chi_i \chi_j^* |i\rangle\langle j| \tag{21}$$

where χ_i are the coordinates of ψ in the computational basis: $|\psi\rangle = \sum_{i=0}^{n-1} \chi_i |i\rangle$. We decompose the system into a u-dimensional subsystem, A, and a v-dimensional subsystem, B, such that $n = uv$, and expand the density matrix ρ in the orthonormal basis $|i\rangle\langle j| = |k\rangle\langle l| \otimes |q\rangle\langle r|$, where each vector is a member of the computational basis in its respective space. The density matrix then takes the form

$$\rho = \sum_{k=0}^{u-1}\sum_{l=0}^{u-1}\sum_{q=0}^{v-1}\sum_{r=0}^{v-1} \chi_{kq}\chi_{lr}^* |k\rangle\langle l| \otimes |q\rangle\langle r| \tag{22}$$

where we have reparameterised the χ_i coordinates as χ_{kq}. We now define the partial traces over the two subsystems as

$$\mathrm{Tr}_A(\rho) = \sum_{t=0}^{u-1} \left(\langle t| \otimes \mathbb{1}_B\right) \rho \left(|t\rangle \otimes \mathbb{1}_B\right) \tag{23}$$

$$\mathrm{Tr}_B(\rho) \;=\; \sum_{t=0}^{v-1} (\mathbb{1}_A \otimes \langle t|)\, \rho\, (\mathbb{1}_A \otimes |t\rangle) \tag{24}$$

where $\mathbb{1}_A$ and $\mathbb{1}_B$ are the identity matrices in the subspaces of A and B, respectively. The reduced density matrix ρ_A, defined by tracing out over the subsystem B, can then be written as

$$\begin{aligned}\rho_A = \mathrm{Tr}_B(\rho) \;&=\; \sum_{t=0}^{v-1}\sum_{k=0}^{u-1}\sum_{l=0}^{u-1}\sum_{q=0}^{v-1}\sum_{r=0}^{v-1} \chi_{kq}\chi^*_{lr}|k\rangle\langle l| \otimes \langle t|q\rangle\langle r|t\rangle \\ &=\; \sum_{t=0}^{v-1}\sum_{k=0}^{u-1}\sum_{l=0}^{u-1} \chi_{kt}\chi^*_{lt}|k\rangle\langle l| \end{aligned} \tag{25}$$

or, in terms of components,

$$(\rho_A)_{kl} = \sum_{t=0}^{v-1} \chi_{kt}\chi^*_{lt} \tag{26}$$

Similarly, the reduced density matrix ρ_B, defined by tracing out over the subsystem A, can be written as

$$(\rho_B)_{qr} = \mathrm{Tr}_A(\rho) = \sum_{t=0}^{u-1} \chi_{tq}\chi^*_{tr} \tag{27}$$

In terms of the reduced density matrices ρ_A and ρ_B of Eqs.(26) and (27), we find that the extremal conditions of Eqs.(19) can be written as

$$\sum_{i=0}^{u-1} (\rho_A)_{ki}\, a_i \;=\; N_A N_B a_k \tag{28}$$

$$\sum_{j=0}^{v-1} (\rho_B)_{mj}\, b_j \;=\; N_A N_B b_m \tag{29}$$

Thus, $N_A N_B$ can be interpreted as the eigenvalues of the reduced density matrices ρ_A and ρ_B, with a_i and b_j being the corresponding eigenvectors. As discussed after Eq. (2), this then provides a geometric interpretation of the Schmidt decomposition of $|\psi\rangle$: the coefficients a_i and b_j used in defining the closest separable state are related to the basis states $|\alpha_k\rangle$ and $|\beta_k\rangle$ of the Schmidt decomposition, with the norm $N_A N_B$ of the closest separable state related to the Schmidt coefficients p_k.

It is interesting to consider the particular case that the n–dimensional space is split into a product of a single qubit space and another space of dimension $u = n/2$. In this case, one of the equations of (20) will become a quadratic equation for the product $N_A N_B$, with solutions

$$N_A N_B \equiv \mu_\pm = \frac{1}{2}\left[1 \pm \sqrt{1-4C}\right], \qquad\qquad C = \sum_{j=1}^{u-1}\sum_{k=0}^{j-1} |\chi_{0j}\chi_{1k} - \chi_{1j}\chi_{0k}|^2 \tag{30}$$

The Schmidt decomposition of Eq. (2) in this case becomes

$$|\psi\rangle = \sqrt{\mu_+}\; |\alpha_+\rangle \otimes |\beta_+\rangle + \sqrt{\mu_-}\; |\alpha_-\rangle \otimes |\beta_-\rangle \tag{31}$$

with $\mu_+ + \mu_- = 1$. Relating the eigenvalues $N_A N_B$ to the cosine of the angle between $|\psi\rangle$ and the closest separable state $|\phi\rangle$ by Eq. (12), and noting that μ_+ is the larger of the two eigenvalues, we find the Schmidt decomposition can be written as

$$|\psi\rangle = \cos\theta_{\max}\; |\alpha_+\rangle \otimes |\beta_+\rangle + \sin\theta_{\max}\; |\alpha_-\rangle \otimes |\beta_-\rangle \tag{32}$$

where $\cos\theta_{\max} \equiv \mu_+$. If $|\psi\rangle$ was separable we would have $\sin^2\theta_{\max} = \mu_- = 0$, and as such this provides a direct connection between the entanglement measures of $1 - \Lambda^2_{\max} = \sin^2\theta_C$ in the geometric approach and the coefficient $\sin^2\theta_{\max}$ in the Schmidt decomposition.

This geometric connection can also be made to the generalization of the Schmidt decomposition for multipartite pure states developed by Partovi [14]. In this approach, one starts with a state $|\psi\rangle$ and decomposes it into two subsystems: one, a qubit space A, and another space $(BC\ldots Z)$ representing the remaining dimensions:

$$|\psi\rangle = \sum_{i_a} \sqrt{p^A_{i_a}} |\psi^A_{i_a}\rangle \otimes |\psi^{BC\ldots Z}_{i_a}\rangle \tag{33}$$

One then decomposes $|\psi^{BC\ldots Z}_{i_a}\rangle$ into two subsystems: another qubit space B, and another space $(CD\ldots Z)$ representing the remaining dimensions:

$$|\psi^{BC\ldots Z}_{i_a}\rangle = \sum_{i_b} \sqrt{p^B_{i_a;i_b}} |\psi^B_{i_a;i_b}\rangle \otimes |\psi^{CD\ldots Z}_{i_a;i_b}\rangle \tag{34}$$

This process is continued until the last two qubit spaces Y and Z are reached, with the result

$$|\psi\rangle = \sum_{i_a i_b \ldots i_{yz}} \sqrt{p^A_{i_a} p^B_{i_a;i_b} \cdots p^{YZ}_{i_a i_b \ldots i_x;i_{yz}}} |\psi^A_{i_a}\rangle \otimes |\psi^B_{i_a;i_b}\rangle \otimes \cdots \otimes |\psi^Y_{i_a i_b \ldots i_x;i_{yz}}\rangle \otimes |\psi^Z_{i_a i_b \ldots i_x;i_{yz}}\rangle \tag{35}$$

There is a direct correspondence between each stage of this series of decompositions and a problem involving the finding of the extremal points of a particular distance in the geometric approach described in the previous section. For example, at the first stage of Eq. (33), we can consider the distance between $|\psi\rangle$ and and a state $|\phi^{A;BC\ldots Z}\rangle = |\phi^A\rangle \otimes |\phi^{BC\ldots Z}\rangle$:

$$D^2_{A;BC\ldots Z} = \langle \psi - \left[\phi^A \otimes \phi^{BC\ldots Z}\right] | \psi - \left[\phi^A \otimes \phi^{BC\ldots Z}\right] \rangle \tag{36}$$

where $|\phi^A\rangle$ is a qubit state and $|\phi^{BC\ldots Z}\rangle$ encompasses the remaining dimensions. Finding the extremal points of this distance will result in a system of (linear) equations, as in Eq. (20), determining the components of the state $|\phi^{A;BC\ldots Z}\rangle$. For the next stage, corresponding to Eq. (34), we can then consider the distance between the state $|\phi^{BC\ldots Z}\rangle$ and a state $|\phi^{B;CD\ldots Z}\rangle = |\phi^B\rangle \otimes |\phi^{CD\ldots Z}\rangle$:

$$D^2_{B;CD\ldots Z} = \langle \phi^{BC\ldots Z} - \left[\phi^B \otimes \phi^{CD\ldots Z}\right] | \phi^{BC\ldots Z} - \left[\phi^B \otimes \phi^{CD\ldots Z}\right] \rangle \tag{37}$$

where $|\phi^B\rangle$ is a qubit state and $|\phi^{CD\ldots Z}\rangle$ encompasses the remaining dimensions. Finding the extremal points of this distance will again result in a system of linear equations determining the components of the state $|\phi^{B;CD\ldots Z}\rangle$. This process may be continued until the last two qubit states $|\phi^Y\rangle$ and $|\phi^Z\rangle$ are reached; at each stage there will be a direct correspondence between the coefficients used in defining the closest separable state to the basis states of the Schmidt decomposition, with the norm of the closest separable state related to the corresponding Schmidt coefficients. At the end, we can then define, in analogy with Eq.(12), the cosine of the critical angle θ_C as

$$\cos\theta_C = \left. \frac{\langle\psi|\phi\rangle}{\sqrt{\langle\phi|\phi\rangle}\sqrt{\langle\psi|\psi\rangle}} \right|_{\text{critical}} \qquad \ldots \quad |\phi\rangle = |\phi^A\rangle \otimes |\phi^B\rangle \otimes \cdots |\phi^Z\rangle \tag{38}$$

and then use as a measure of entanglement $\sin^2\theta_C = 1 - \cos^2\theta_C$. Although this procedure has the advantage compared to the approach of Section II of resulting in a series of linear equations to solve, compared to the non–linear equations of Eq. (9), the disadvantage is that the final result depends on the order that the series of decompositions is made: the sequence $ABC\ldots YZ$ described above will differ from the sequence $BC\ldots ZA$. As such, an approach such as that of Ref. [14] of a minimization over all permutations of the possible orders of the decompositions must be done.

IV. CONCLUSIONS

We have considered a generalization of the usual geometric measure of entanglement of pure states using the distance to the nearest unnormalized product state. Although this doesn't lead to any computational advantages, as the resulting equations determining the measure are still non–linear in general, this does afford an interpretation of the standard entanglement measure $1 - \Lambda^2_{\max}$ as the distance to the closest separable state. This also provides a relationship between the the norm and components of the closest separable state and the coefficients and basis states of the Schmidt decomposition of the state $|\psi\rangle$.

Acknowledgments

This work was supported by the Natural Sciences and Engineering Research Council of Canada.

[1] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
[2] Ryszard Horodecki, Pawel Horodecki, Michal Horodecki, and Karol Horodecki, arXiv:quant-ph/0702225 (submitted to Rev. Mod. Phys.).
[3] Martin B. Plenio and Shashank Virmani, Quant. Inf. Comp. **7**, 1 (2007).
[4] Karol Zyczkowski and Ingemar Bengtsson, arXiv:quant-ph/0606228v1.
[5] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Phys. Rev. Lett. **78**, 2275 (1997).
[6] A. Shimony, Ann. N. Y. Acad. Sci. **755**, 675 (1995).
[7] H. Barnum and N. Linden, J. Phys. **A34**, 6787 (2001).
[8] Tzu-Chieh Wei and Paul M. Goldbart, Phys. Rev. **A68**, 042307 (2003).
[9] Ya Cao and An Min Wang, arXiv:quant-ph/0701099v2.
[10] Tzu-Chieh Wei and Paul M. Goldbart, arXiv:quant-ph/0303079v1
[11] Tsubasa Ichikawa, Izumi Tsutsui, and Taksu Cheon, arXiv:quant-ph/0702167.
[12] Jon Magne Leinaas, Jan Myrheim, and Eirik Ovrum, arXiv:quant-ph/0605079.
[13] A. Yu. Bogdanov, Yu. I. Bogdanov, and K. A. Valiev, arXiv:quant-ph/0512062.
[14] M. Hossein Partovi, Phys. Rev. Lett. **92**, 077904 (2004).
[15] Ashish V. Thapliyal, Phys. Rev. **A59**, 3336 (1999).